

Eigenvalues and Expansion of Regular Graphs

Nabil Kahale*

Abstract

The spectral method is the best currently known technique to prove lower bounds on expansion. Ramanujan graphs, which have asymptotically optimal second eigenvalue, are the best known explicit expanders. The spectral method yielded a lower bound of $k/4$ on the expansion of linear sized subsets of k -regular Ramanujan graphs. We improve the lower bound on the expansion of Ramanujan graphs to approximately $k/2$. Moreover, we construct a family of k -regular graphs with asymptotically optimal second eigenvalue and linear expansion equal to $k/2$. This shows that $k/2$ is the best bound one can obtain using the second eigenvalue method. We also show an upper bound of roughly $1 + \sqrt{k-1}$ on the average degree of linear-sized induced subgraphs of Ramanujan graphs. This compares positively with the classical bound $2\sqrt{k-1}$. As a byproduct, we obtain improved results on random walks on expanders and construct selection networks (resp. extrovert graphs) of smaller size (resp. degree) than was previously known.

Categories and Subject Descriptors: F.2.2 [**Analysis of Algorithms and Problems Complexity**]: Nonnumerical Algorithms and Problems; G.2.2 [**Discrete Mathematics**]: Graph Theory
General Terms: Algorithms, Theory

Additional Key Words and Phrases: Eigenvalues, Expander graphs, Induced subgraphs, Load balancing, Ramanujan graphs, Random walks, Selection networks.

1 Introduction

Expander graphs are widely used in Theoretical Computer Science, in areas ranging from parallel computation [1, 7, 21, 28, 34] to complexity theory and cryptography [2, 8, 16, 35]. Given an

*XEROX Palo Alto Research Center, 3333 Coyote Hill Road, CA 94304. This work was done mostly when the author was at the Massachusetts Institute of Technology, and partly at DIMACS. This work was partially supported by the Defense Advanced Research Projects Agency under Contracts N00014-92-J-1799 and N00014-91-J-1698, the Air Force under Contract F49620-92-J-0125, and the Army under Contract DAAL-03-86-K-0171. This paper was based on “Better Expansion for Ramanujan graphs”, by Nabil Kahale, which appeared in the *32nd Annual Symposium on Foundations of Computer Science*, San Juan, Puerto Rico, October 1–4, 1991; pp. 398–404. ©IEEE, and on “On the Second Eigenvalue and Linear Expansion of Regular Graphs”, by Nabil Kahale, which appeared in the *33rd Annual Symposium on Foundations of Computer Science*, Pittsburgh, Pennsylvania, October 24–27, 1992; pp. 296–303. ©IEEE. An updated version of the second paper appeared in *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, Volume 10, 1993; pp. 49–62. ©American Mathematical Society.

undirected k -regular graph $G = (V, E)$ and a subset X of V , the expansion of X is defined to be the ratio $|N(X)|/|X|$, where $N(X) = \{w \in V : \exists v \in X, (v, w) \in E\}$ is the set of neighbors of X . An (α, β, k, n) -expander is a k -regular graph on n nodes such that every subset of size at most αn has expansion at least β .

It is known that random regular graphs are good expanders. In particular, for any $\beta < k - 1$, there exists a constant α such that, with high probability, all the subsets of a random k -regular graph of size at most αn have expansion at least β . The explicit construction of expander graphs is much more difficult, however. The first explicit construction of an infinite family of expanders was discovered by Margulis [25] in 1973, and improved in [15, 5, 17].

The best currently known method to calculate lower bounds on the expansion in polynomial time relies on analyzing the second eigenvalue of the graph. Since the adjacency matrix A is symmetric, all its eigenvalues are real and will be denoted by $\lambda_0 \geq \lambda_1 \geq \dots \geq \lambda_{n-1}$. We have $\lambda_0 = k$, and $\lambda = \max(\lambda_1, |\lambda_{n-1}|) \leq k$. Tanner [33] proved that for any subset X of V ,

$$|N(X)| \geq \frac{k^2|X|}{\lambda^2 + (k^2 - \lambda^2)|X|/n}. \quad (1)$$

Therefore, in order to get high expansion, we need λ to be as small as possible. However, for any sequence $G_{n,k}$ of k -regular graphs on n vertices, $\liminf \lambda(G_{n,k}) \geq 2\sqrt{k-1}$ as n goes to infinity [3, 23, 27]. Therefore, the best expansion coefficient we can obtain by applying Tanner's result is approximately $k/4$. This bound is achieved by Ramanujan graphs, which have been explicitly constructed [23, 26] for many pairs (k, n) . By definition, a Ramanujan graph is a connected k -regular graph whose eigenvalues $\neq \pm k$ are at most $2\sqrt{k-1}$ in absolute value. The relationship between the eigenvalues of the adjacency matrix and the expansion coefficient has also been investigated in [3, 5, 6, 12], but the bound they get, when applied to non-bipartite Ramanujan graphs and for sufficiently large k , is no better than Tanner's bound. Other results about expanders are contained in [9, 22, 30].

Some applications, such as the construction of non-blocking networks in [7], required an expansion greater than $k/2$ for linear-sized subsets. Indeed, if the expansion of a subset X is greater than $k/2$, a constant fraction its nodes have *unique neighbors*, i.e. neighbors adjacent to only one node in X . This allows the construction of a matching between X and $N(X)$ in a logarithmic number of steps and using only *local computations*. Recently, Pippenger [29] showed that weak expanders are sufficient in applications where an expansion greater than $k/2$ was required.

We define the linear expansion of a family of graphs G_n on n vertices to be the best lower bound on the expansion of subsets of size up to αn , where α is an arbitrary small positive constant, that is

$$\sup_{\alpha > 0} \inf_n \inf_X \frac{|N(X)|}{|X|},$$

where X ranges over the subsets of G_n of size at most αn . We show that if (G_n) is a family of k -regular graphs whose second largest eigenvalue is upper bounded by $\tilde{\lambda}$, the linear expansion of (G_n) is at least $(k/2) \left(1 - \sqrt{1 - (4k-4)/\tilde{\lambda}^2}\right)$. In particular, the expansion of linear-sized subsets of Ramanujan graphs is lower bounded by a factor arbitrary close to $k/2$. On the other hand, for any integer k such that $k-1$ is a prime congruent to 1 modulo 4, and for any function m of n such that $m = o(n)$, we explicitly construct an infinite family of k -regular graphs G_n on n vertices such that $\lambda(G_n) \leq (2 + o(1))\sqrt{k-1}$ and G_n contains a subset of size $2m$ with expansion $k/2$. Since such a family has asymptotically optimal second eigenvalue, this shows that $k/2$ is essentially the best lower

bound on the linear expansion one can obtain by the second eigenvalue method. The techniques used in this construction can be applied to prove tightness of relationships between eigenvalues and diameter [20]. Our results provide an efficient way to test that the expansion of linear sized subsets of random graphs is at least $k/2 - O(k^{3/4} \log^{1/2} k)$. We also show that the average degree of the induced subgraphs on linear-sized subsets of a k -regular graph G is upper bounded by a factor arbitrary close to $1 + \tilde{\lambda}/2 + \sqrt{\tilde{\lambda}^2/4 - (k-1)}$, where $\tilde{\lambda} = \max(\lambda, 2\sqrt{k-1})$. This bound is equal to $1 + \sqrt{k-1}$ in the case of Ramanujan graphs, improving upon the previously known bound [4] of $2\sqrt{k-1}$.

Sections 3–5 contain our main results. In Section 6, we apply our techniques to obtain improved results on random walks on expanders. Random walks are often used in complexity theory and cryptography, and our bound improves upon previous results in [2, 16]. Applications to selection networks and extrovert graphs are described in Section 7. We conclude with some remarks in Section 8.

Some of the results in this paper have appeared in an extended abstract form in [18, 19], and in a more detailed form in [20].

2 Notation, definitions, and background

Throughout the paper, $G = (V, E)$ will denote an undirected graph on a set V of vertices. Let $L^2(V)$ denote the set of real valued functions on V and $L_0^2(V) = \{f \in L^2(V); \sum_{v \in V} f(v) = 0\}$. As usual, we define the scalar product of two vectors f and g of $L^2(V)$ by

$$f \cdot g = \sum_{v \in V} f(v)g(v),$$

and the euclidean norm of a vector f by $\|f\| = \sqrt{f \cdot f}$. We denote the adjacency matrix of G by A_G , or simply by A if there is no risk of confusion. The matrix A is the 0-1 $n \times n$ matrix whose (i, j) entry is equal to 1 if and only $(i, j) \in E$. It defines a linear operator in $L^2(V)$ that maps every vector $f \in L^2(V)$ to the vector Af defined by

$$(Af)(v) = \sum_{(v,w) \in E} f(w). \quad (2)$$

This operator is selfadjoint since $\forall f, g \in L^2(V)$,

$$(Af) \cdot g = f \cdot (Ag) = \sum_{(v,w) \in E} f(v)g(w). \quad (3)$$

For any matrix or operator M with real eigenvalues, we denote by $\lambda_i(M)$ the i -th largest eigenvalue of M , $\lambda_i(A_G)$ by $\lambda_i(G)$, and $\max(\lambda_1(G), |\lambda_{n-1}(G)|)$ by $\lambda(G)$. For any subset W of V , we denote by χ_W the characteristic vector of W : $\chi_W(v) = 1$, if $v \in W$, and 0 otherwise. We denote the adjacency matrix of the graph induced on W by A_W , the real number $\lambda_i(A_W)$ by $\lambda_i(W)$, and the set of nodes at distance at most l from W by $B_l(W)$. For the rest of this section, we assume that G is k -regular.

Fact 1 [24, 32] *If B is a selfadjoint operator in a vector space L , then*

$$\lambda_0(B) = \max_{g \in L - \{0\}} \frac{g \cdot Bg}{\|g\|^2}.$$

■

Clearly, the vector χ_V is an eigenvector of A with eigenvalue k . The vector space $L_0^2(V)$ is invariant under A , and the eigenvalues of the restriction of A to $L_0^2(V)$ are $\lambda_1(G), \dots, \lambda_{n-1}(G)$. Therefore,

Fact 2 For any $g \in L_0^2(V)$, we have $g \cdot Ag \leq \lambda_1(G) \|g\|^2$. ■

For two column vectors g and h , we say that $g \leq h$ if every coordinate of g is at most its corresponding coordinate in h .

Fact 3 [31, page 28, ex. 1.12] If a real symmetric matrix has only nonnegative entries, its largest eigenvalue is nonnegative and has a corresponding eigenvector with nonnegative components. This eigenvalue is largest in absolute value. ■

Fact 4 If a real symmetric matrix B has only nonnegative entries, and s is a vector with positive components such that $Bs \leq \gamma s$, then the largest eigenvalue of B is at most γ . This property still holds if only the off-diagonal entries of B are assumed to be nonnegative [14]. ■

Given a graph H , the *cover graph* H' of H is the graph defined on $V' = V \times \{0, 1\}$ and where $((u, l), (v, m)) \in V' \times V'$ is an edge if and only $(u, v) \in E$ and $l \neq m$.

Fact 5 The eigenvalues of H' consist of the eigenvalues of H and their negated values. ■

3 Main lemma

In this section, we prove the main lemma (Theorem 1) that we will use later to derive lower bounds on the expansion.

Lemma 1 If $G = (V, E)$ is k -regular on n vertices, then for any $f \in L^2(V)$, we have

$$f \cdot Af \leq \lambda_1(G) \|f\|^2 + \frac{k - \lambda_1(G)}{n} \left(\sum_{v \in V} f(v) \right)^2.$$

Proof We decompose f as the sum of a constant vector and a vector in $L_0^2(V)$. Let $\bar{f} = (f \cdot \chi_V / n) \chi_V$ be the orthogonal projection of f on the subspace spanned by the constant vector χ_V . Then $f_0 = f - \bar{f}$ is the orthogonal projection of f on $L_0^2(V)$. By linearity, $Af = A\bar{f} + Af_0 = k\bar{f} + Af_0$, and so $f \cdot Af = k \|\bar{f}\|^2 + f_0 \cdot Af_0$ since $\bar{f} \cdot Af_0 = A\bar{f} \cdot f_0 = k\bar{f} \cdot f_0 = 0$. By Fact 2 and the Pythagorean theorem, we have

$$f_0 \cdot Af_0 \leq \lambda_1(G) \|f_0\|^2 = \lambda_1(G) (\|f\|^2 - \|\bar{f}\|^2).$$

Therefore, $f \cdot Af \leq \lambda_1(G) \|f\|^2 + (k - \lambda_1(G)) \|\bar{f}\|^2$. We conclude the proof by noting that

$$\|\bar{f}\|^2 = \left(\sum_{v \in V} f(v) \right)^2 / n.$$

■

Lemma 2 For any subset W of a k -regular graph G , we have $\lambda_0(W) \leq \lambda_1(G) + (k - \lambda_1(G)) |W| / n$.

Proof Let g be any element of $L^2(W)$. Consider the vector $f \in L^2(V)$ that coincides with g on $L^2(W)$ and is null on $V - W$. By Eq. 3, we see that $g \cdot A_W g = f \cdot A_G f$. By applying Lemma 1 to f , we have

$$\begin{aligned} g \cdot A_W g &\leq \lambda_1(G) \|g\|^2 + \frac{k - \lambda_1(G)}{n} \left(\sum_{v \in W} g(v) \right)^2 \\ &\leq \left(\lambda_1(G) + \frac{k - \lambda_1(G)}{n} |W| \right) \|g\|^2. \end{aligned}$$

The second equation follows from the Cauchy-Schwartz inequality. We conclude using Fact 1. ■

A similar relation [20, Section 4.1] holds between $\lambda_i(W)$ and $\lambda_{i+1}(G)$, for $1 \leq i \leq |W| - 1$. Lemma 2 already gives a restriction on the structure of induced subgraphs. For example, since the average degree of a graph is upper bounded by its largest eigenvalue, it implies that the average degree of the induced subgraph on W is at most $\lambda_1(G) + (k - \lambda_1(G))|W|/n$, which is roughly $\lambda_1(G)$ for small linear-sized subsets. To obtain a stronger restriction on the induced subgraph on a linear-sized subset X , we will apply Lemma 2 to the set $B_l(X)$. We start by comparing the largest eigenvalue of the subgraph induced on $B_l(X)$ to the matrix of a weighted graph associated with X .

Let Δ_X be the diagonal matrix indexed by the vertices of X and whose entry (v, v) is equal to the degree of v in the subgraph induced on X . We will simply denote by $\Delta_X(v)$ the diagonal entry $\Delta_X(v, v)$. For $\theta' > 0$ and integer l , let

$$M_X^{\theta', l} = A_X + \frac{1}{\sqrt{k-1}} \frac{\sinh(l\theta')}{\sinh((l+1)\theta')} (kI - \Delta_X).$$

The matrix $M_X^{\theta', l}$ can be regarded as the matrix of the weighted graph that is induced on X and has in addition a loop of weight $(k-1)^{-1/2} (\sinh(l\theta')/\sinh((l+1)\theta'))(k - \Delta_X(v))$ on each node v of X .

Lemma 3 *Suppose $G = (V, E)$ is k -regular. Let l be a positive integer, X a non-empty subset of V , θ' a positive real number, and $\lambda' = 2\sqrt{k-1} \cosh \theta'$. If $\lambda_0(B_l(X)) \leq \lambda'$, then $\lambda_0(M_X^{\theta', l}) \leq \lambda'$.*

Proof Let $W = B_l(X)$. The idea behind the proof is as follows. Let $f \in L^2(X)$ be an eigenvector of $M_X^{\theta', l}$ corresponding to its largest eigenvalue. We will extend f to W so that it becomes roughly an eigenvector of A_W . If the largest eigenvalue of $M_X^{\theta', l}$ were too big, we would get a large eigenvalue for A_W , contradicting the fact that $\lambda_0(W) \leq \lambda'$.

We define the sequence r_i as follows:

$$r_i = \frac{\sinh((l+1-i)\theta')(k-1)^{-i/2}}{\sinh((l+1)\theta')}.$$

The sequence r_i is strictly positive and decreasing for $0 \leq i \leq l$, and

$$\lambda' r_i = r_{i-1} + (k-1)r_{i+1} \tag{4}$$

$$\lambda' r_l = r_{l-1}. \tag{5}$$

By Fact 3, we can assume that all the entries of f are nonnegative. We extend the vector f to W by setting

$$f(v) = \max_{u \in X} (f(u)r_{d(v,u)}),$$

for $v \in W - X$, where $d(v, u)$ denote the distance in G between v and u . Note that f is nonnegative on W since every node in W is at distance at most l from some node in X .

Claim 1 For any $v \in W - X$, we have $(A_W f)(v) \geq \lambda' f(v)$.

Proof Let $u \in X$ be such that $f(v) = f(u)r_{d(v,u)}$. Note that $i = d(v, u)$ can be assumed to be at most l since $r_j \leq 0$ for $j \geq l + 1$. Let v_1 be the first node on a shortest path from v to u . Since the distance between v_1 and u is $i - 1$, we have $v_1 \in W$ and

$$f(v_1) \geq f(u)r_{i-1}. \quad (6)$$

We now distinguish two cases:

- **Case 1.** $i = l$. Combining Eqs. 6 and 5, we get in this case

$$(A_W f)(v) \geq f(v_1) \geq f(u)r_{i-1} = \lambda' f(u)r_i = \lambda' f(v),$$

as required.

- **Case 2.** $i < l$. In this case, the k neighbors of v are at distance at most $i + 1 \leq l$ from u . By monotonicity of the sequence r , it follows that the value of f on each of these neighbors is at least $f(u)r_{i+1}$. Using again Eq. 6, we have

$$\begin{aligned} (A_W f)(v) &\geq f(v_1) + (k - 1)f(u)r_{i+1} \\ &\geq f(u)(r_{i-1} + (k - 1)r_{i+1}) \\ &= \lambda' f(u)r_i \\ &= \lambda' f(v). \end{aligned}$$

■

Claim 2 For any $v \in X$, we have $(A_W f)(v) \geq \lambda_0(M_X^{\theta', l})f(v)$.

Proof For any $v \in X$, the value of f on each of the $k - \Delta_X(v)$ neighbors of v in $W - X$ is at least $f(v)r(1)$. Therefore,

$$(A_W f)(v) \geq (A_X f)(v) + (k - \Delta_X(v))f(v)r(1) = (M_X^{\theta', l} f)(v) = \lambda_0(M_X^{\theta', l})f(v).$$

■

Since $\lambda_0(W) \leq \lambda'$, we have $(A_W f) \cdot f \leq \lambda' \|f\|^2$, by Fact 1. Assume for contradiction that $\lambda' < \lambda_0(M_X^{\theta', l})$. By Claim 1 and Claim 2, this implies that $(A_W f) \cdot f > \lambda' \|f\|^2$, leading to a contradiction. ■

For $\theta \geq 0$, define

$$M_X^\theta = A_X + \frac{1}{\sqrt{k-1}e^\theta}(kI - \Delta_X).$$

The matrix M_X^θ can be regarded as the matrix of the weighted graph that is induced on X and has in addition a loop of weight $(k - 1)^{-1/2}e^{-\theta}(k - \Delta_X(v))$ on each node v of X .

Theorem 1 Suppose $G = (V, E)$ is k -regular, and let $\tilde{\lambda} = \max(\lambda_1(G), 2\sqrt{k-1}) = 2\sqrt{k-1} \cosh \theta$, where $\theta \geq 0$. For any non-empty subset X of V of size at most $k^{-1/\epsilon}|V|$, we have

$$\lambda_0(M_X^\theta) \leq \tilde{\lambda}(1 + O(\epsilon)),$$

where the constant behind the O is a small absolute constant.

Proof Let $l = \lfloor 1/2\epsilon \rfloor$ and let W be the set of nodes at distance at most l from X . A simple calculation shows that $|W| \leq 3k^l |X| \leq 3k^{-1/(2\epsilon)} n$. It follows from Lemma 2 that $\lambda_0(W) \leq \lambda'$, where $\lambda' = \tilde{\lambda} + 3k^{1-1/(2\epsilon)} = 2\sqrt{k-1} \cosh \theta'$, with $\theta' > \theta \geq 0$. A straightforward calculation shows that $\lambda' = \tilde{\lambda}(1+O(\epsilon))$ and $\cosh \theta' - \cosh \theta = O(k^{1/2-1/(2\epsilon)}) = O(\epsilon^2)$. We will use the following inequalities to show that the matrix $M_X^{\theta',l}$ is approximated by M_X^θ .

Claim 3 For $x \geq y \geq 0$, we have $(x-y)^2 \leq 2(\cosh x - \cosh y)$.

Proof This follows immediately from Taylor's expansion formula. ■

Claim 4

$$\frac{l}{l+1} e^{-\theta'} \leq \frac{\sinh(l\theta')}{\sinh((l+1)\theta')} \leq e^{-\theta'}.$$

Proof Since $\sinh((l+1)\theta') \geq (l+1)\sinh \theta'$, we have

$$e^{-\theta'} - \frac{\sinh(l\theta')}{\sinh((l+1)\theta')} = \frac{e^{-l\theta'} - e^{-(l+2)\theta'}}{e^{(l+1)\theta'} - e^{-(l+1)\theta'}} = e^{-(l+1)\theta'} \frac{\sinh \theta'}{\sinh((l+1)\theta')} \leq \frac{e^{-\theta'}}{l+1}.$$

■

It follows from Claim 3 that $\theta' - \theta = O(\epsilon)$. On the other hand, Claim 4 implies that

$$\frac{\sinh(l\theta')}{\sinh((l+1)\theta')} = e^{-\theta'}(1 + O(1/l)) = e^{-\theta}(1 + O(\epsilon)).$$

Therefore, all entries of the diagonal matrix $M_X^\theta - M_X^{\theta',l}$ are $O(\sqrt{k-1}\epsilon)$, and so its largest eigenvalue is $O(\sqrt{k-1}\epsilon)$. But, as a consequence of Fact 1, the function that associates to a symmetric matrix its largest eigenvalue is subadditive. Therefore,

$$\begin{aligned} \lambda_0(M_X^\theta) &\leq \lambda_0(M_X^{\theta',l}) + O(\sqrt{k-1}\epsilon) \\ &\leq \lambda' + O(\tilde{\lambda}\epsilon) \\ &= \tilde{\lambda}(1 + O(\epsilon)), \end{aligned}$$

where the second inequality follows from Lemma 3. ■

Remark 1 The only place where we used in the proof the fact that $\lambda_1(G)$ is the second eigenvalue of G was in conjunction with Lemma 2 to upper bound $\lambda_0(W)$. In particular, if λ^* is a real number such that for any subset W of V , we have $\lambda_0(W) \leq \lambda^* + 2k|W|/|V|$, then Theorem 1 remains valid if $\lambda_1(G)$ is replaced by λ^* .

4 Lower bounds on the expansion and on the average degree

We will derive lower bounds on the expansion by applying Theorem 1 to the union of X and $N(X)$, after reducing to the case where the graph is bipartite and X is on one side of the partition. The idea is that if the expansion of X is small, a node in $N(X)$ will be adjacent to many nodes in X , in average. This implies that the largest eigenvalue of the weighted matrix associated to the subgraph induced on $X \cup N(X)$ is large, contradicting with Theorem 1. We also use Theorem 1 to derive upper bounds on the average degree of induced subgraphs.

Theorem 2 *If $G = (V, E)$ is k -regular and $\tilde{\lambda} = \max(\lambda_1(G), 2\sqrt{k-1})$, then for any non-empty subset X of V of size at most $k^{-1/\epsilon}|V|$,*

$$\frac{|N(X)|}{|X|} \geq \frac{k}{2} \left(1 - \sqrt{1 - \frac{4k-4}{\tilde{\lambda}^2}} \right) (1 - O(\epsilon)),$$

where the constant behind the O is a small absolute constant.

Proof We first show how to reduce the problem to the case where the graph G is bipartite and X is on one side of the partition. Consider the cover graph G' of G , as defined in Section 3. We show that Remark 1 applies to the graph G' and $\lambda^* = \lambda_1(G)$. Indeed, let W be a subset of V' , $W_p \subseteq V$ the set of nodes u of V such that $(u, 0) \in W$ or $(u, 1) \in W$, and $W^* = W_p \times \{0, 1\}$. Since the largest eigenvalue of a graph is no less than the largest eigenvalue of any induced subgraph [10, page 156], $\lambda_0(W) \leq \lambda_0(W^*)$. (We remind the reader that $\lambda_0(W)$ (resp. $\lambda_0(W^*)$) is the largest eigenvalue of the subgraph of G' induced on W (resp. W^*), and $\lambda_0(W_p)$ is the largest eigenvalue of the subgraph of G induced on W_p .) On the other hand, it follows from Fact 5 and Fact 3 that $\lambda_0(W^*) = \lambda_0(W_p)$. Using Lemma 2, we get

$$\lambda_0(W) \leq \lambda_0(W_p) \leq \lambda_1(G) + k \frac{|W_p|}{|V|} = \lambda_1(G) + 2k \frac{|W|}{|V'|},$$

as required by Remark 1. Note that we cannot apply directly Theorem 1 because, by Fact 5, we have $\lambda_1(G') = \lambda(G)$, which may be different from $\lambda_1(G)$.

Let Y be the subset of V' equal to $X \times \{0\}$. Denote the adjacency matrix of G' by A' , and the set of neighbors of Y in G' by $N'(Y)$. Let $\tilde{\lambda} = 2\sqrt{k-1} \cosh \theta$, with $\theta \geq 0$. By applying Remark 1 to the graph G' , the set of vertices $Y \cup N'(Y)$, and $\epsilon' = 2\epsilon$, we see that the largest eigenvalue of the matrix $M' = M'_{Y \cup N'(Y)}$ is at most $\tilde{\lambda}(1 + O(\epsilon))$. Now, consider the function $f \in L^2(Y \cup N'(Y))$ defined by $f = k\chi_Y + \tilde{\lambda}\chi_{N'(Y)}$. By Fact 1,

$$M'f \cdot f \leq \tilde{\lambda}(1 + O(\epsilon))\|f\|^2. \quad (7)$$

The left-hand side is the sum of two terms. The first is equal to $A'f \cdot f$, and the second corresponds to the weighted self-loops. By Eq. 3, we have $A'f \cdot f = 2\tilde{\lambda}k^2|Y|$. On the other hand, since the loops have no weight on Y and have average weight $e^{-\theta}(k-1)^{-1/2}(k-k|Y|/|N'(Y)|)$ on $N'(Y)$, the second term is equal to $e^{-\theta}(k-1)^{-1/2}(k|N'(Y)| - k|Y|)\tilde{\lambda}^2$. Thus Eq. 7 reduces to

$$2\tilde{\lambda}k^2|Y| + \frac{\tilde{\lambda}^2}{\sqrt{k-1}e^\theta}k(|N'(Y)| - |Y|) \leq \tilde{\lambda}(1 + O(\epsilon))(k^2|Y| + \tilde{\lambda}^2|N'(Y)|).$$

By replacing $|Y|$ by $|X|$, $|N'(Y)|$ by $|N(X)|$, we get after simplifications

$$k|X|(k - 2e^{-\theta} \cosh \theta) \leq |N(X)|(\tilde{\lambda}^2 - 2ke^{-\theta} \cosh \theta)(1 + O(\epsilon)). \quad (8)$$

Noting that $\tilde{\lambda}^2 - 2ke^{-\theta} \cosh \theta = 2(k e^\theta - 2 \cosh \theta) \cosh \theta$, Eq. 8 reduces to:

$$\frac{|N(X)|}{|X|} \geq \frac{k}{2e^\theta \cosh \theta} (1 - O(\epsilon)).$$

We conclude the proof using the formula

$$\frac{1}{e^\theta \cosh \theta} = 1 - \sqrt{1 - \frac{1}{\cosh^2 \theta}} = 1 - \sqrt{1 - \frac{4k-4}{\tilde{\lambda}^2}}.$$

■

Theorem 3 *If $G = (V, E)$ is k -regular and $\tilde{\lambda} = \max(\lambda_1(G), 2\sqrt{k-1})$, then for any non-empty subset X of V of size at most $k^{-1/\epsilon}|V|$, the average degree σ of the subgraph of G induced on X is at most*

$$\left(1 + \frac{\tilde{\lambda}}{2} + \sqrt{\frac{\tilde{\lambda}^2}{4} - (k-1)}\right)(1 + O(\epsilon)),$$

where the constant behind the O is a small absolute constant.

Proof We use the same notations as in Theorem 1. As noted before, the matrix M_X^θ can be regarded as the matrix of the weighted graph on X that is induced on X and has in addition a loop of weight $(k-1)^{-1/2}e^{-\theta}(k - \Delta_X(v))$ on each node v of X . By Fact 1, we have $\chi_X \cdot M_X^\theta \chi_X \leq \lambda_0(M_X^\theta)|X|$, which translates into

$$\sigma + \frac{k - \sigma}{\sqrt{k-1}e^\theta} \leq 2\sqrt{k-1}(1 + O(\epsilon)) \cosh \theta.$$

This implies

$$\begin{aligned} \sigma &\leq \frac{2(k-1)e^\theta \cosh \theta - k}{\sqrt{k-1}e^\theta - 1}(1 + O(\epsilon)) \\ &= (\sqrt{k-1}e^\theta + 1)(1 + O(\epsilon)). \end{aligned}$$

We conclude by noting that $e^\theta = \cosh \theta + \sqrt{\cosh^2 \theta - 1}$. ■

5 A family of almost Ramanujan graphs with expansion $k/2$

In this section, we construct explicitly a family of k -regular graphs G_n containing subsets of sublinear size having expansion $k/2$, and such that $\lambda(G_n) = (2 + o(1))\sqrt{k-1}$. For this, we need the following lemma.

Lemma 4 *Consider a graph on a vertex set W , a subset X of W , a positive integer h , and $s \in L^2(W)$. Let X_i be the set of nodes at distance i from X . Assume the following conditions hold:*

1. *For $h-1 \leq i, j \leq h$, all nodes in X_i have the same number of neighbors in X_j .*
2. *The vector s is constant on X_{h-1} and on X_h .*
3. *s has positive components and $As \leq \mu s$ on $B_{h-1}(X)$, where μ is a positive real number.*

Then for any $g \in L^2(W)$ such that $|Ag(u)| = \mu|g(u)|$ for $u \in B_{h-1}(X)$, we have

$$\frac{\sum_{v \in X_h} g(v)^2}{\sum_{v \in X_h} s(v)^2} \geq \frac{\sum_{v \in X_{h-1}} g(v)^2}{\sum_{v \in X_{h-1}} s(v)^2}.$$

Proof Let P, P_{h-1} and P_h be the projections on the sets $B_{h-1}(X), X_{h-1}$ and X_h respectively. We need to show that $\|P_h g\|^2 / \|P_h s\|^2 \geq \|P_{h-1} g\|^2 / \|P_{h-1} s\|^2$. Let $A_h = (P + P_h)A(P + P_h)$. The operator A_h corresponds in some sense to the adjacency matrix of the subgraph induced on $B_h(X)$, but it acts on $L^2(W)$. By the conditions of the lemma, there exist positive coefficients α, β and γ

such that $P_h A_h s = \gamma P_h s$ and $A_h P_h s = \alpha P_h s + \beta P_{h-1} s$. By hypothesis, we have $A_h s \leq \mu P s + \gamma P_h s$. Premultiplying both sides of this equation by P and A_h yields successively

$$\begin{aligned} P A_h s &\leq \mu P s \\ &= \mu s - \mu P_h s, \\ A_h P A_h s &\leq \mu A_h s - \mu(\alpha P_h s + \beta P_{h-1} s) \\ &\leq \mu^2 P s + \mu(\gamma - \alpha) P_h s - \mu\beta P_{h-1} s. \end{aligned}$$

But the matrix $A_h P A_h - \mu^2 P - \mu(\gamma - \alpha) P_h + \mu\beta P_{h-1}$ has only nonnegative entries off its diagonal, and so its largest eigenvalue is 0 since s is positive (Fact 4.) The quadratic form associated to this matrix is therefore negative semi-definite (Fact 1), and so

$$A_h P A_h g \cdot g \leq \mu^2 P g \cdot g + \mu(\gamma - \alpha) P_h g \cdot g - \mu\beta P_{h-1} g \cdot g. \quad (9)$$

Since both A_h and P are selfadjoint and since $P^2 = P$, the left-hand side of Eq. 9 is equal to $\|P A_h g\|^2$. We can rewrite Eq. 9 as follows:

$$\|P A_h g\|^2 \leq \mu^2 \|P g\|^2 + \mu(\gamma - \alpha) \|P_h g\|^2 - \mu\beta \|P_{h-1} g\|^2.$$

But $\|P A_h g\| = \mu \|P g\|$ by hypothesis, and so

$$(\gamma - \alpha) \|P_h g\|^2 \geq \beta \|P_{h-1} g\|^2. \quad (10)$$

On the other hand, since A_h and P_h are selfadjoint, we have $A_h P_h s \cdot s = P_h A_h s \cdot s$, and so $\alpha \|P_h s\|^2 + \beta \|P_{h-1} s\|^2 = \gamma \|P_h s\|^2$. Comparing this with Eq. 10 concludes the proof. \blacksquare

Theorem 4 *For any integer k such that $k - 1$ is prime, we can explicitly construct an infinite family of k -regular graphs G_n on n vertices whose linear expansion is $k/2$ and such that $\lambda_1(G_n) \leq 2\sqrt{k-1}(1 + 2\log^2 \log n / \log_k^2 n)$.*

Proof We construct the family (G_n) by altering the known constructions of explicit Ramanujan graphs, so that the expansion of (G_n) is $k/2$. From [23] and [26], we know that we can explicitly construct an infinite family of bipartite Ramanujan graphs (F_n) on n vertices whose girth $c(F_n)$ is $(4/3 + o(1)) \log_{k-1} n$. Let $F_n = (V, E)$ be an element of the family, $u \in V$ a vertex of F_n and $l = \lfloor c(F_n)/2 \rfloor - 2$. Let u_1, \dots, u_k be the neighbors of u and let v_1, \dots, v_k be k vertices at distance two from u such that $(u_i, v_i) \in E$. The subgraph of F_n induced on $B_{l+1}(\{u\})$ is a tree since it contains no cycles. Let u' and v' be two elements not belonging to V . Consider the k -regular graph $G_{n+2} = (V', E')$, where $V' = V \cup \{u', v'\}$ and $E' = E \cup \bigcup_{i=1}^k \{(u', u_i), (u_i, u'), (v', v_i), (v_i, v')\} - \bigcup_{i=1}^k \{(u_i, v_i), (v_i, u_i)\}$. Fig. 1 shows the graph G_{n+2} in the neighborhood of u in the case $k = 3$. For shorthand, we denote A_{F_n} by A , $A_{G_{n+2}}$ by A' and $\lambda_1(A')$ by λ' . We need to show that $\lambda' \leq (2 + o(1))\sqrt{k-1}$. Assume that $\lambda' > 2\sqrt{k-1}$ (otherwise we are done), and let $\lambda' = 2\sqrt{k-1} \cosh \theta'$, with $\theta' > 0$. Let $g \in L_0^2(V')$ be an eigenvector corresponding to λ' .

We outline informally the basic ideas of the proof. Roughly speaking, we will show that the values that g takes on the nodes u', v', u_i, v_i are small compared to $\|g\|$. This implies that g is close in l_2 -norm to its restriction f on V . Lemma 1 then implies that $f \cdot A f \leq (2 + o(1))\sqrt{k-1} \|g\|^2$. But, since $g(u')$, $g(v')$, $g(u_i)$ and $g(v_i)$ are small, the scalar product $f \cdot A f$ is close to $g \cdot A' g = \lambda' \|g\|^2$, and so $\lambda' \leq (2 + o(1))\sqrt{k-1}$.




Figure 1: The graph G_{n+2} in the neighborhood of u in the case $k = 3$. The dotted edges are those belonging to $E - E'$.

Since u and u' have the same neighbors in G_{n+2} and $\lambda' \neq 0$, we have $g(u) = g(u')$. By Eq. 3, we have

$$\begin{aligned} \lambda' \|g\|^2 &= g \cdot A'g & (11) \\ &= f \cdot Af - 2 \sum_{i=1}^k g(u_i)g(v_i) + 2 \sum_{i=1}^k g(u')g(u_i) + 2 \sum_{i=1}^k g(v')g(v_i). \end{aligned}$$

We upper bound $-2g(u_i)g(v_i)$ by $g(u_i)^2 + g(v_i)^2$. On the other hand, the equality $(A'g)(u') = \lambda'g(u')$ implies that

$$\sum_{i=1}^k g(u')g(u_i) = \frac{1}{\lambda'} \left(\sum_{i=1}^k g(u_i) \right)^2 \leq \frac{k}{\lambda'} \sum_{i=1}^k g(u_i)^2.$$

A similar relation holds for v' . Combining this with Eq. 11, we get

$$\lambda' \|g\|^2 \leq f \cdot Af + \left(1 + \frac{2k}{\lambda'}\right) \sum_{i=1}^k (g(u_i)^2 + g(v_i)^2). \quad (12)$$

We use Lemma 1 to bound the term $f \cdot Af$. Note that $\sum_{w \in V} f(w) = -g(u') - g(v')$ since $g \in L_0^2(V')$, and so

$$\begin{aligned} f \cdot Af &\leq \lambda_1(A) \|f\|^2 + \frac{k}{n} (g(u') + g(v'))^2 \\ &\leq 2\sqrt{k-1} (\|g\|^2 - g(u')^2 - g(v')^2) + \frac{2k}{n} (g(u')^2 + g(v')^2) \\ &\leq 2\sqrt{k-1} \|g\|^2, \end{aligned}$$

for sufficiently large n . Combining this with Eq. 12 and noting that $1 + 2k/\lambda' \leq 4\sqrt{k-1}$, we obtain

$$\lambda' \|g\|^2 \leq 2\sqrt{k-1} (\|g\|^2 + 2 \sum_{i=1}^k (g(u_i)^2 + g(v_i)^2)). \quad (13)$$

We now compare $g(u_i)$ and $g(v_i)$ to $\|g\|$. Let s be the function on V' defined by $s(u) = s(u') = k$ and, for $v \in V' - \{u, u'\}$ at distance i from u , $s(v) = 2(k-1)^{1-i/2} \cosh i\theta'$. An easy calculation shows that the function s verifies the conditions of Lemma 4 for the graph G' , with $X = \{u, u'\}$, $\mu = \lambda'$, and for any integer h between 1 and $l+1$. Hence

$$\begin{aligned} \|g\|^2 &\geq \sum_{v \in X_{l+1}} g(v)^2 \\ &\geq \frac{\sum_{v \in X_{l+1}} s(v)^2}{\sum_{v \in X_1} s(v)^2} \sum_{v \in X_1} g(v)^2 \\ &= \frac{k-2}{k-1} \frac{\cosh^2(l+1)\theta'}{\cosh^2 \theta'} \sum_{i=1}^k g(u_i)^2 \\ &\geq \frac{1}{2} \cosh^2(l\theta') \sum_{i=1}^k g(u_i)^2. \end{aligned}$$

Similarly, $\|g\|^2 \geq \frac{1}{2} \cosh^2(l\theta') \sum_{i=1}^k g(v_i)^2$. Combining this with Eq. 13, we get

$$\cosh \theta' \leq 1 + \frac{8}{\cosh^2 l\theta'}. \quad (14)$$

Solving Eq. 14 yields $\theta' \leq (\log l)/l$ for sufficiently large n , and so

$$\lambda' = 2\sqrt{k-1} \left(1 + \frac{\theta'^2}{2} (1 + o(1))\right) \leq 2\sqrt{k-1} \left(1 + 2 \frac{\log^2 \log n}{\log_k^2 n}\right).$$

Theorem 2 implies that the linear expansion of the family (G_n) is at least $k/2$. Since the subset $\{u, u'\}$ has k neighbors, this bound is tight. ■

If $k-1$ is a prime congruent to 1 modulo 4, we know from [23] that there exists an infinite family of non-bipartite k -regular Ramanujan graphs with girth at least $(2/3 + o(1)) \log_{k-1} n$. By repeating the construction in Theorem 4, we obtain k -regular graphs whose second largest eigenvalue in absolute value is $(2 + o(1))\sqrt{k-1}$ and linear expansion $k/2$. Moreover, by adjoining nodes at regions of the graph at sufficiently large distance from each other, we can construct for any $m = m(n) = o(n)$ a family of k -regular graphs whose second largest eigenvalue in absolute value is $(2 + o(1))\sqrt{k-1}$ and containing a subset of size $2m$ with expansion $k/2$. This can be shown by a proof similar to Theorem 4.

6 Random walks

We show that the probability that a walk stays inside a given set has an exponential decay in the length of the walk. Our bound improves upon previous results in [2, 16], and is shown to be optimal for many values of the parameters.

Corollary 1 *If $G = (V, E)$ is k -regular and W a subset of V , the fraction of walks in G of length l whose all vertices belong to W is at most $\mu (\alpha + \mu - \alpha\mu)^l$, where $\alpha = \lambda_1/k$ and $\mu = |W|/n$.*

Proof The number of walks of length l in W is equal to the sum of entries of A_W^l , and so the fraction of walks of length l in W is equal to $((A_W)^l \chi_W \cdot \chi_W) / (k^l n)$. On the other hand, since

$\lambda_0(A_W)$ is the largest eigenvalue of A_W in absolute value, the largest eigenvalue of A_W^l is $(\lambda_0(A_W))^l$. Using Lemma 2 and Fact 1, we conclude that

$$(A_W^l \chi_W) \cdot \chi_W \leq (\lambda_0(A_W))^l \|\chi_W\|^2 \leq \mu n (\lambda_1 + k\mu - \lambda_1\mu)^l,$$

as desired. ■

Define the density of a subset of vertices to be the ratio of its size to the total number of vertices. Corollary 1 is optimal in the following sense: for any rational $\alpha \in [1/2, 1]$ and any rational $\mu \in [0, 1]$, there exists an arbitrary large k -regular graph G and a subset W of G of density μ such that $\lambda_1(G) = \alpha k$ and, for any integer l , the fraction of walks in W of length l is equal to $\mu (\alpha + \mu - \alpha\mu)^l$. Indeed, let $G = K_{a+2} \times K_b$, where $b = a(1 - \alpha)/\alpha$ and $K_{a+2} \times K_b$ is the graph on $V = \{1, \dots, a+2\} \times \{1, \dots, b\}$, with $((i, j), (i', j')) \in E$ if and only if $i = i'$ XOR $j = j'$. The graph G is regular of degree $a + b$; its eigenvalues are $a + b, a, b - 2$ and -2 , and $\lambda_1(G) = a = \alpha(a + b)$. Let $W = \{1, \dots, a+2\} \times \{1, \dots, \mu b\}$. The set W has density μ in V , and the fraction of walks in W of length l is $\mu(a + \mu b)^l / (a + b)^l$, which is equal to the value given by Corollary 1.

7 Other applications

We list three applications of Theorems 2, 3.

1. **Random regular graphs.** It was shown in [13] that, if k is even, then for a random k -regular graphs G , we have $\lambda_1(G) \leq 2\sqrt{k-1} + O(\log k)$ with high probability. Using Theorem 2, we deduce that for a random regular graph, we can prove with high probability in polynomial time that linear sized subsets (of density at most $k^{-1/\epsilon}$, where $\epsilon = k^{-1/4}$) have expansion at least

$$\frac{k}{2} \left(1 - \sqrt{1 - \frac{4k-4}{(2\sqrt{k-1} + O(\log k))^2}} \right) (1 + O(k^{-1/4})) = \frac{k}{2} - O(k^{3/4} \log^{1/2} k).$$

2. **Selection networks.** We can use Theorem 2 to build explicit selection networks of small size. A selection network is a network of comparators that classifies a set of n numbers, where n is even, into two subsets of $n/2$ numbers such that any element in the first subset is smaller than any element in the second subset. In [28], a probabilistic construction of a selection network is given using an asymptotic upper bound of $2n \log_2 n$ comparators. Also, an upper bound slightly less than $6n \log_2 n$ is shown by a deterministic construction. Using Theorem 2, we can construct selection networks of asymptotic size $(3 + o(1))n \log_2 n$. Indeed, it is shown in [28] how to construct selection networks of size $(2 + o(1))n \log_2 n$ from expanders of degree 4 having linear expansion at least 3. The construction can be easily generalized to build selection networks of size $k(1/2 + o(1))n \log_2 n$ from expanders of degree k having linear expansion at least 3. Theorem 2 then shows that we can build explicit selection networks of size $(3 + o(1))n \log_2 n$ using 6-regular Ramanujan graphs.
3. **Extrovert graphs.** Given a graph $G = (V, E)$ and a subset X of V , an element of X is said to be extrovert if at least half of its neighbors are outside X . A family of graphs is called extrovert if all linear-sized subsets contain a constant fraction of extrovert nodes. Such graphs have been used [11] to solve the token distribution problem. Theorem 3 shows that

the average degree of the nodes of a linear-sized induced subgraph of a k -regular Ramanujan graph is upper bounded by roughly $1 + \sqrt{k-1}$, which is less than $k/2$ for $k \geq 7$. This shows that Ramanujan graphs of degree at least 7 are extrovert graphs. Classical results [4] require the degree to be at least 15.

8 Concluding remarks and further work

1. Let H be a graph of maximum degree at most k , and $\bar{\lambda}$ a real number no smaller than $2\sqrt{k-1}$. Theorem 1 implies that, if there exists an infinite family G_n of k -regular graphs containing H as an induced subgraph and such that $\lambda(G_n) \leq (1 + o(1))\bar{\lambda}$, then $\lambda_0(M_H^\theta) \leq \bar{\lambda}$. (M_H^θ can be defined similarly to M_X^θ .) If $k-1$ is a prime congruent to 1 modulo 4, this condition can be shown to be sufficient [20].
2. It is still an open question whether there exists a family of Ramanujan graphs with linear expansion at most $k/2$.
3. It would be interesting to calculate the exact value of the linear expansion of the known explicit constructions of Ramanujan graphs [23, 26]. Theorem 2 shows that it is at least $k/2$. On the other hand, an easy combinatorial argument shows that the linear expansion of any family of k -regular graphs is at most $k-1$. Besides being Ramanujan, the graphs constructed in [23, 26] have other interesting combinatorial properties. For example, they are Cayley graphs and have high girth, unlike the graphs that we constructed in Section 5. This leads us to conjecture that their linear expansion is strictly greater than $k/2$. Any explicit construction of k -regular graphs with provable linear expansion strictly greater than $k/2$ would also be interesting.

References

- [1] M. Ajtai, J. Komlós, and E. Szemerédi. Sorting in $c \log n$ parallel steps. *Combinatorica*, 3:1–19, 1983.
- [2] M. Ajtai, J. Komlós, and E. Szemerédi. Deterministic simulation in logspace. In *Proceedings of the 19th Annual ACM Symposium on Theory of Computing*, pages 132–140. ACM Press, 1987.
- [3] N. Alon. Eigenvalues and expanders. *Combinatorica*, 6(2):83–96, 1986.
- [4] N. Alon and F. R. K. Chung. Explicit construction of linear sized tolerant networks. *Discrete Mathematics*, 72:15–19, 1988.
- [5] N. Alon, Z. Galil, and V. D. Milman. Better expanders and superconcentrators. *J. Algorithms*, 8:337–347, 1987.
- [6] N. Alon and V. D. Milman. λ_1 , isoperimetric inequalities for graphs and superconcentrators. *Journal of Combinatorial Theory Series B*, 38:73–88, 1985.
- [7] S. Arora, T. Leighton, and B. Maggs. On-line algorithms for path selection in a non-blocking network. In *Proceedings of the 22nd Annual ACM Symposium on Theory of Computing*, pages 149–158. ACM Press, 1990.
- [8] M. Bellare, O. Goldreich, and S. Goldwasser. Randomness in interactive proofs. In *Proceedings of the 31st Annual Symposium on Foundations of Computer Science*, pages 563–572, 1990.

- [9] F. Bien. Constructions of telephone networks by group representations. *Notices Amer. Math. Soc.*, 36(1):5–22, 1989.
- [10] B. Bollobás. *Graph Theory*. Springer-Verlag, New York, 1990.
- [11] A. Z. Broder, A. M. Frieze, E. Shamir, and E. Upfal. Near-perfect token distribution. In *19th International Colloquium on Automata, Languages, and Programming*, pages 308–317. Springer Verlag, W. Berlin, 1992.
- [12] M. W. Buck. Expanders and diffusers. *SIAM J. Algebraic Discrete Methods*, 7(2):282–304, 1986.
- [13] J. Friedman. On the second eigenvalue and random walks in random regular graphs. *Combinatorica*, 11(4):331–362, 1991.
- [14] J. Friedman. Some geometric aspects of graphs and their eigenfunctions. *Duke Math. J.*, 69(3):487–525, 1993.
- [15] O. Gabber and Z. Galil. Explicit constructions of linear sized superconcentrators. *J. Comput. System Sci.*, 22(3):407–420, 1981.
- [16] O. Goldreich, R. Impagliazzo, L. Levin, R. Venkatesan, and D. Zuckerman. Security preserving amplification of hardness. In *Proceedings of the 31st Annual Symposium on Foundations of Computer Science*, pages 318–326. IEEE Computer Society Press, 1990.
- [17] S. Jimbo and A. Maruoka. Expanders obtained from affine transformations. *Combinatorica*, 7(4):343–355, 1987.
- [18] N. Kahale. Better expansion for Ramanujan graphs. In *Proceedings of the 32nd Annual Symposium on Foundations of Computer Science*, pages 296–303. IEEE Computer Society Press, 1991.
- [19] N. Kahale. On the second eigenvalue and linear expansion of regular graphs. In *Proceedings of the 33rd Annual Symposium on Foundations of Computer Science*, pages 398–404. IEEE Computer Society Press, October 1992. An updated version appeared in “DIMACS Series in Discrete Mathematics and Theoretical Computer Science, Volume 10, 1993, 49–62”.
- [20] N. Kahale. *Expander Graphs*. PhD thesis, Massachusetts Institute of Technology, September 1993. MIT Laboratory for Computer Science, Technical Report MIT/LCS/TR-591.
- [21] T. Leighton and B. Maggs. Fast algorithms for routing around faults on multibutterflies. In *Proceedings of the 30th Annual Symposium on Foundations of Computer Science*, 1989.
- [22] A. Lubotzky. *Discrete groups, expanding graphs and invariant measures*. Manuscript.
- [23] A. Lubotzky, R. Phillips, and P. Sarnak. Ramanujan graphs. *Combinatorica*, 8(3):261–277, 1988.
- [24] M. Marcus and H. Minc. *A Survey of Matrix Theory and Matrix Inequalities*. Dover, 1992.
- [25] G. A. Margulis. Explicit constructions of concentrators. *Problemy Peredači Informacii*, 9(4):71–80, 1973.
- [26] G. A. Margulis. Explicit group-theoretical constructions of combinatorial schemes and their applications to the design of expanders and concentrators. *Problemy Peredači Informacii*, 24(1):51–60, 1988.

- [27] A. Nilli. On the second eigenvalue of a graph. *Discrete Mathematics*, 91:207–210, 1991.
- [28] N. Pippenger. Selection networks. *SIAM J. of Comput.*, 20(5):878–887, 1991.
- [29] N. Pippenger. Self-routing superconcentrators. In *Proceedings of the 25th Annual ACM Symposium on Theory of Computing*, 1993.
- [30] P. Sarnak. *Some applications of modular forms*. Cambridge University Press, 1990.
- [31] E. Seneta. *Non-negative matrices and Markov Chains*. Springer-Verlag, 1981.
- [32] G. Strang. *Linear Algebra and its Applications*. Harcourt Brace Jovanovich, San Diego, 1988.
- [33] R. M. Tanner. Explicit construction of concentrators from generalized n -gons. *SIAM J. Algebraic Discrete Methods*, 5(3):287–294, 1984.
- [34] E. Upfal. An $O(\log n)$ deterministic packet routing scheme. In *Proceedings of the 21st Annual ACM Symposium on Theory of Computing*, pages 241–250. ACM Press, 1989.
- [35] L. Valiant. Graph theoretic properties in computational complexity. *J. Comput. System Sci.*, 13:278–285, 1976.